



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

On the Structure of Forms, and the Algebraical Theory of n -Lines.

BY O. E. GLENN.

Factorable ternary quantics, representing n -lines, have an invariant theory which is in some respects analogous to binary invariant theory. The theory of degenerate forms lends considerable aid also to the study of non-factorable forms. In this paper we propose to develop the theory of factorable forms considerably farther than was done in a paper on the subject published by the present writer in Vol. XXXII, No. 1, of this JOURNAL,* and to make a series of applications of this theory. Sections 1, 2, 3, 4 are devoted to ternary factor theorems. In § 5 an introduction to an invariant theory of n -lines from the standpoint of matrices is given. Resultant and discriminant matrices of forms representing n -lines are constructed. Section 6 contains a theory of rational partial fractions from the point of view of the Aronhold operator.

§ 1. *Multiple Linear Factors.*

The number of terms in an elementary symmetric function of m groups of any p homogeneous variables is equal to the number of distinct permutations of the variables occurring in any one term, when the subscripts are removed. Thus if the groups are

$$\begin{aligned} &P_1(q_1, r_1, s_1), \\ &P_2(q_2, r_2, s_2), \\ &\dots\dots\dots, \\ &P_m(q_m, r_m, s_m), \end{aligned}$$

we are led by a simple proof to the relation

$$\begin{aligned} &\sum \frac{\partial}{\partial r_1} (\sum q_1 q_2 \dots q_r r_{r+1} r_{r+2} \dots r_{m-k} s_{m-k+1} \dots s_m) s_1 \\ &= (k+1) \sum q_1 q_2 \dots q_r r_{r+1} r_{r+2} \dots r_{m-k-1} s_{m-k} \dots s_m. \end{aligned} \quad (1)$$

* "The Theory of Degenerate Algebraical Curves and Surfaces." The paper contains a short bibliography of the subject, to which may be added, BRIOSCHI: "Sulla Condizioni, etc.," *Annali di Matematica*, Ser. 12, Vol. VII; THAEER: "Ueber die Zerlegbarkeit einer ebenen Linie, u. s. w.," *Math. Annalen*, Vol. XIV; BES: "Décomposition de la forme ternaire du troisième degré," *Math. Annalen*, Vol. LIX (1904).

Hence, when the ternary form

$$f_{3m} = a_{0m} y^m + \sum_{r=0}^{m-1} (a_{m-r} x^{m-r} + a_{m-r-1} x^{m-r-1} + \dots + a_{0r}) y^r,$$

which we write under the non-homogeneous notation

$$f_{3m} = x^m \phi_{0y/x} + x^{m-1} \phi_{1y/x} + \dots + x \phi_{m-1y/x} + \phi_m,$$

is decomposable into m linear factors, its coefficients are connected with the coefficients of these linear forms, as

$$\rho_i = q_i y + r_i x + s_i \quad (i = 1, 2, \dots, m),$$

by the following relations, among others:

$$\sum \frac{\partial a_{m-r-k} r}{\partial r_1} s_1 = (k+1) a_{m-r-k-1} r \begin{pmatrix} r=0, 1, \dots, m-1 \\ k=0, 1, \dots, m-1 \\ \pi q_i = a_{0m} = 1 \end{pmatrix}. \quad (2)$$

Assume that r_1 is a root of multiplicity α_1 of

$$\phi_{0-r} \equiv r^m - a_{1m-1} r^{m-1} + a_{2m-2} r^{m-2} - \dots + (-1)^m a_{m0} = 0.$$

Then evidently r_1 is a root of multiplicity $\alpha_1 - \kappa$ of

$$\phi_{\kappa-r} = 0 \quad (\kappa = 0, 1, 2, \dots, \overline{\alpha_1 - 1}).$$

Now, when the coefficients a of $\phi_{\kappa-r}$ are expressed in terms of the variables r_i, s_i , we write

$$\phi_{\kappa} = \phi_{\kappa-r}(a) = I_{\kappa}(r_i, s_i; r) = I_{\kappa},$$

and then $I_{\kappa}(r_i, s_i; r_1) \equiv 0$. By replacing r_i, r in I_{κ} by $r_i + \lambda s_i, r + \lambda s$ respectively, expanding by Taylor's theorem and using (2), we get

$$\begin{aligned} & \sum_{h=0}^{\infty} \frac{\lambda^h}{h!} \left(\sum \frac{\partial}{\partial r_1} s_1 + \frac{\partial}{\partial r} s \right)^h I_{\kappa} \\ &= \sum_{h=0}^{\infty} \frac{\lambda^h}{h!} \left(\sum \frac{\partial}{\partial a_{m-r-k} r} (k+1) a_{m-r-k-1} r + \frac{\partial}{\partial r} s \right)^h \phi_{\kappa} \\ &= \sum_{h=0}^{\infty} \frac{\lambda^h}{h!} \sum_{i=0}^h \binom{h}{i} s^{h-i} \frac{\partial^{h-i} \Delta_k^i \phi_{\kappa}}{\partial r^{h-i}} \\ &= \sum_{h=0}^{\infty} \frac{\lambda^h}{h!} \sum_{i=0}^h (-1)^i \binom{h}{i} (x+1)(x+2)\dots(x+i) s^{h-i} \frac{\partial^{h-i} \phi_{\kappa+i}}{\partial r^{h-i}}, \end{aligned} \quad (3)$$

where

$$\Delta_k = \sum_k \left(a_{m-k-10} \frac{\partial}{\partial a_{m-k0}} + a_{m-k-21} \frac{\partial}{\partial a_{m-k-11}} + \dots + a_{0m-k-1} \frac{\partial}{\partial a_{1m-k-1}} \right) (k+1).$$

The first non-vanishing term of this series, when $r = r_1$, is the term where $h = \alpha_1 - \kappa$. Placing $\kappa = 0$, we have from this term the result that the roots s_i corresponding to the α_1 equal r_i are the α_1 roots of the equation

$$\frac{\partial^{\alpha_1} \phi_{0-r_1}}{\partial r_1^{\alpha_1}} s^{\alpha_1} - \alpha_1 \frac{\partial^{\alpha_1-1} \phi_{1-r_1}}{\partial r_1^{\alpha_1-1}} s^{\alpha_1-1} + \alpha_1 (\alpha_1 - 1) \frac{\partial^{\alpha_1-2} \phi_{2-r_1}}{\partial r_1^{\alpha_1-2}} s^{\alpha_1-2} - \dots + (-1)^{\alpha_1} \alpha_1 \phi_{\alpha_1-r_1} = 0. \quad (4)$$

Similar results hold for a root r_i of ϕ_{0-r} of multiplicity α_i ($i = 1, 2, \dots, t$). That is, we have for the homogeneous

$$f_{3m} = x_2^m \phi_{0, x_1/x_2} + x_2^{m-1} x_3 \phi_{1, x_1/x_2} + x_2^{m-2} x_3^2 \phi_{2, x_1/x_2} + \dots + \phi_{m, x_1/x_2} x_3^m :$$

THEOREM 1. *The linearly factorable homogeneous ternary form f_{3m} , whose leading binary ϕ_{0-r} has r_i as a root of multiplicity α_i ($i = 1, 2, \dots, t$; $\alpha_1 + \alpha_2 + \dots + \alpha_t = m$), can be factored into factors of the respective orders $\alpha_1, \alpha_2, \dots, \alpha_t$, which are rational and integral in the coefficients of the form f_{3m} itself on the one hand, and in the quantities r_i on the other, linear in the coefficients; according to the formula*

$$f_{3m} = \prod_{i=1}^t \left[\frac{\partial^{\alpha_i} \phi_{0-r_i}}{\partial r_i^{\alpha_i}} x^{\alpha_i} - \alpha_i \frac{\partial^{\alpha_i-1} \phi_{1-r_i}}{\partial r_i^{\alpha_i-1}} x^{\alpha_i-1} y_i + \alpha_i (\alpha_i - 1) \frac{\partial^{\alpha_i-2} \phi_{2-r_i}}{\partial r_i^{\alpha_i-2}} x^{\alpha_i-2} y_i^2 + \dots + (-1)^{\alpha_i} \alpha_i \phi_{\alpha_i-r_i} y_i^{\alpha_i} \right] (x_1 + r_i x_2/x_3 = x/y_i). \quad (5)$$

As an application of this theorem consider the problem of resolving the factorable ternary quartic f_{34} , where

$$x_2^4 \phi_{0, x_1/x_2} = a_{400} x_1^4 + a_{310} x_1^3 x_2 + a_{220} x_1^2 x_2^2 + a_{130} x_1 x_2^3 + a_{040} x_2^4, \text{ etc.,}$$

and

$$\begin{aligned} a_{400} &= 1, & a_{301} &= a + 3, & a_{202} &= 3a - 4, & a_{103} &= -4a - 12, & a_{004} &= -12a, \\ a_{310} &= 9, & a_{211} &= 7a + 19, & a_{112} &= 13a - 22, & a_{013} &= -14a - 24, \\ a_{220} &= 30, & a_{121} &= 16a + 40, & a_{022} &= 14a - 28, \\ a_{130} &= 44, & a_{031} &= 12a + 28, \\ a_{040} &= 24. \end{aligned}$$

We find the roots of $\phi_{0, x_1/x_2} = 0$ to be $+r_1 = +2$, $+r_2 = 3$; $\alpha_1 = 3$, $\alpha_2 = 1$. Equation (4) and the corresponding one for s_2 , α_2 are then computed. They are

$$s^3 - (a + 1)s^2 + (a - 6)s + 6a = 0, \quad s - 2 = 0.$$

Their roots are $s_1 = -2$, $s_2 = 3$, $s_3 = a$; $s_4 = 2$. Hence,

$$f_{34} = (x_1 + 2x_2 - 2x_3)(x_1 + 2x_2 + 3x_3)(x_1 + 2x_2 + ax_3)(x_1 + 3x_2 + 2x_3).$$

§ 2. *Extension of Method for Quaternary Forms.*

For elementary symmetric functions of m groups of four variables we have

$$\begin{aligned} \Sigma \frac{\partial}{\partial r_1} (\Sigma q_1 q_2 \cdots q_{m-j-k-l} r_{m-j-k-l+1} r_{j-k-l+2} \cdots r_{m-j-k} s_{m-j-k+1} \cdots s_{m-j} t_{m-j+1} \cdots t_m) t_j \\ = (j+1) \Sigma q_1 q_2 \cdots q_{m-j-k-l} r_{m-j-k-l+1} r_{m-j-k-l+2} \cdots r_{m-j-k-1} s_{m-j-k} \\ \cdots s_{m-j-1} t_{m-j} \cdots t_m, \\ \Sigma \frac{\partial}{\partial r_1} (\Sigma q_1 q_2 \cdots q_{m-j-k-l} r_{m-j-k-l+1} r_{m-j-k-l+2} \cdots r_{m-j-k} s_{m-j-k+1} \cdots s_{m-j} t_{m-j+1} \cdots t_m) s_1 \\ = (k+1) \Sigma q_1 q_2 \cdots q_{m-j-k-l} r_{m-j-k-l+1} r_{m-j-k-l+2} \cdots r_{m-j-k+1} s_{m-j-k} \\ \cdots s_{m-j} t_{m-j+1} \cdots t_m, \end{aligned}$$

and with

$$\begin{aligned} f_{4m} &= \sum_{l=0}^m \left[\sum_{k=0}^{m-l} (a_{lm-k-lk} x^l + a_{l-1m-k-lk} x^{l-1} + \cdots + a_{0m-k-lk} z^k) y^{m-k-l} \right. \\ &\quad \left. = \prod_{i=1}^m (q_i y + r_i x + s_i z + t_i), \right. \end{aligned}$$

there results

$$\left. \begin{aligned} \Sigma \frac{\partial a_{lm-j-k-lk}}{\partial r_1} t_1 &= (j+1) a_{l-1m-j-k-lk}, \\ \Sigma \frac{\partial a_{lm-j-k-lk}}{\partial r_1} s_1 &= (k+1) a_{l-1m-j-k-lk+1} \end{aligned} \right\} \quad (6)$$

$(j=0, 1, \dots, m-1; k=0, 1, \dots, m-1; l=1, 2, \dots, m).$

The terms of a quaternary form may be arranged, under a homogeneous notation, as follows:

$$\begin{aligned} f_{4m} &= x_2^m \phi_{0 \ x_1/x_2} + x_2^{m-1} x_3 \phi_{1 \ x_1/x_2}^{(1)} + x_2^{m-2} x_3^2 \phi_{2 \ x_1/x_2}^{(1)} + \cdots + x_3^m \phi_{m \ x_1/x_2}^{(1)} \\ &\quad + x_2^{m-1} x_4 \phi_{1 \ x_1/x_2}^{(2)} + x_2^{m-2} x_4^2 \phi_{2 \ x_1/x_2}^{(2)} + \cdots + x_4^m \phi_{m \ x_1/x_2}^{(2)} \\ &\quad + \Psi_{4m}, \end{aligned}$$

where

$$\begin{aligned} x_2^{m-k} \phi_{k \ x_1/x_2}^{(j)} &= a_{m-k-0 \ 0 \dots 0 \ k \ 0 \dots 0}^{\overbrace{j-1} \quad \overbrace{2-j}} x_1^{m-k} + a_{m-k-11 \ 0 \dots 0 \ k \ 0 \dots 0}^{\overbrace{j-1} \quad \overbrace{2-j}} x_1^{m-k-1} x_2 + \dots \\ &\quad + a_{m-k-h \ 0 \dots 0 \ k \ 0 \dots 0}^{\overbrace{j-1} \quad \overbrace{2-j}} x_1^{m-k-h} x_2^h + \cdots + a_{0 \ m-k \ 0 \dots 0 \ k \ 0 \dots 0}^{\overbrace{j-1} \quad \overbrace{2-j}} x_2^{m-k}. \end{aligned}$$

Hence, by means of (6) and the methods of the previous section, we have, after writing the results in homogeneous form, with

$$f_{4m} = \prod_{i=1}^m (x_1 + r_i x_2 + r_{3i} x_3 + r_{4i} x_4) \quad (a_{m000} = 1),$$

THEOREM 2: If r_i is a root of $\phi_{0-r} = 0$ of multiplicity α_i ($i = 1, 2, \dots, t$; $\alpha_1 + \alpha_2 + \dots + \alpha_t = m$) of a quaternary form f_{4m} , then the coefficients r_{j+2i} of its linear factors are the roots of the equation

$$\prod_{i=1}^t \left[\frac{\partial^{a_i} \phi_{0-r_i}^{(j)}}{\partial r_i^{a_i}} r_{j+2}^{a_i} - \alpha_i \frac{\partial^{a_i-1} \phi_{1-r_i}^{(j)}}{\partial r_i^{a_i-1}} r_{j+2}^{a_i-1} + \alpha_i (\alpha_i - 1) \frac{\partial^{a_i-2} \phi_{2-r_i}^{(j)}}{\partial r_i^{a_i-2}} r_{j+2}^{a_i-2} \right. \\ \left. + \dots + (-1)^i [\alpha_i \phi_{a_i-r_i}^{(j)}] \right] = 0 \quad (j = 1, 2; \phi_{0-r}^{(1)} = \phi_{0-r}^{(2)} = \phi_{0-r}). \quad (7)$$

§ 3. Multiple Roots of Functions $\Phi_\xi^{(m)}$, $\Psi_\eta^{(m)}$.

Let the m roots of the equation

$$f(x) = a_0 x^m + a_1 x^{m-1} + \dots + a_m = 0$$

be x_1, x_2, \dots, x_m . Represent the totality of the sums of these roots taken two at a time by ξ_i ($i = 1, 2, \dots, (m) = \frac{1}{2} m(m-1)$), and the totality of the products of the roots taken in pairs by $-\eta_j$ ($j = 1, 2, \dots, (m)$). Let the equations having the sets of quantities ξ_i, η_j for roots be respectively

$$\Phi_\xi^{(m)} = \phi_0 \xi^{(m)} + \phi_1 \xi^{(m)-1} + \dots + \phi_{(m)} = 0, \\ \Psi_\eta^{(m)} = \psi_0 \eta^{(m)} + \psi_1 \eta^{(m)-1} + \dots + \psi_{(m)} = 0.$$

Then, for convenience of statement in this paper, let the following terminology be adopted: Let $x^2 - \xi_1 x - \eta_1$ be a quadratic factor of $f(x)$. This factor will be said to be of sum-multiplicity α if $f(x)$ has exactly α quadratic factors in which the coefficient of x is the same number, $-\xi_1$. The factor will be said to be of product-multiplicity β if just β of the factors of $f(x)$ have the same absolute term $-\eta_1$. Again, the factor will be called of multiplicity γ if it is repeated as a whole just γ times in $f(x)$. Evidently if the factor is of multiplicity γ , its sum-multiplicity or its product-multiplicity, or both, may be equal to γ or greater than γ .

Let $x^2 - \xi_1 x - \eta_1$ be a factor of $f(x)$ of multiplicity α . Suppose its sum-multiplicity and product-multiplicity are both α also. Then x_1, x_2 ($x_1 + x_2 = \xi_1$) are roots of multiplicity α of $f(x)$.

Let the roots of $\Phi_\xi^{(m)} = 0$ be arranged in a triangular array as follows:

$$\begin{array}{ccccccc} x_1 + x_2, & x_1 + x_3, & x_1 + x_4, & x_1 + x_5, & \dots, & & \\ & x_2 + x_3, & x_2 + x_4, & x_2 + x_5, & \dots, & & \\ & & x_3 + x_4, & x_3 + x_5, & \dots, & & \\ & & & x_4 + x_5, & \dots, & & \end{array}$$

Then it is evident that ξ_1 is, *formally*, a root of $\Phi_\xi^{(m)} = 0$ of multiplicity

$$1 + 3 + 5 + \dots + (2\alpha - 1) = \alpha^2.$$

Similarly, η_1 is a root of $\Psi_\eta^{(m)} = 0$ of multiplicity α^2 . More generally, if we assume

$$\left. \begin{aligned} \xi_1 = \xi_2 = \dots = \xi_{a_1+1} = \dots = \xi_{a_1+a_2} = \xi_{a_1+a_2+1} = \dots = \xi_{a_1+a_2+a_3} = \dots = \xi_{\sum_{i=1}^k a_i}, \\ \eta_1 = \eta_2 = \dots = \eta_{a_1} \neq \eta_{a_1+1} = \dots = \eta_{a_1+a_2} \neq \eta_{a_1+a_2+1} = \dots = \eta_{a_1+a_2+a_3} \neq \dots = \eta_{\sum_{i=1}^k a_i}, \end{aligned} \right\} \quad (8)$$

where $\sum_{i=1}^k \alpha_i = \alpha$, it results that ξ_1 is a root of $\Phi_\xi^{(m)} = 0$ of multiplicity

$$\alpha_1^2 + \alpha_2^2 + \dots + \alpha_k^2,$$

whereas $\eta_{a_1+a_2+\dots+a_k}$ is, *formally*, a root of $\Psi_\eta^{(m)}$ of multiplicity α_h^2 ($h=1, 2, \dots, k$).

A dual result of the same nature is obtained by interchanging the rôles of ξ_i, η_i in this paragraph.

§ 4. *Multiple Quadratic Factors.*

Assume that the general ternary form f_{3m} is decomposable into $\frac{1}{2}m = g$ quadratic factors (m even):

$$\tau_i = n_i x^2 + o_i x y + p_i y^2 + q_i x + r_i y + s_i \quad (i = 1, 2, \dots, g).$$

Then

$$\prod_{i=1}^g (p_i y^2 + o_i y x + n_i x^2) = x^m \Phi_{0y/x}.$$

The coefficients of the quadratic forms τ_i and those of f_{3m} itself are connected by the relations

$$\left(\sum \frac{\partial}{\partial n_1} q_1 + \sum \frac{\partial}{\partial o_1} r_1 + 2 \sum \frac{\partial}{\partial q_1} s_1 \right) a_{m-k-r-r} = (k+1) a_{m-k-r-1r} \quad (9)$$

$$(k = 0, 1, 2, \dots, m-1).$$

Consider now the case where $r^2 + o_1 r + n_1$ is a factor of multiplicity 1, and of product-multiplicity 1 and sum-multiplicity α of Φ_{0r} . Then o_1 is a root of multiplicity α of $\Phi_{-o}^{(m)}$, whereas n_1 is a simple root of $\Psi_{-n}^{(m)}$. (We have taken $p_1 = p_2 = \dots = a_{0m} = 1$.)

$$\begin{aligned} \text{Let} \quad \Phi &= \Phi_{-o}^{(m)}(\alpha) \equiv I(n_i, o_i, q_i, \dots; o) = I, \\ \Psi &= \Psi_{-n}^{(m)}(\alpha) \equiv J(n_i, o_i, q_i, \dots; n) = J. \end{aligned}$$

$$\text{Then} \quad I(n_i, o_i, q_i, \dots; o_1) \equiv J(n_i, o_i, q_i, \dots; n_1) \equiv 0.$$

In $\Phi = I$ let us replace n_i, o_i, q_i by $n_i + \lambda q_i, o_i + \lambda r_i, q_i + 2\lambda s_i$ respectively, and o by $o + \lambda r$. Then Taylor's expansion becomes, by virtue of (9),

$$\begin{aligned} & \sum_{j=0}^{\infty} \frac{\lambda^j}{j!} \left(\sum \frac{\partial}{\partial n_1} q_1 + \sum \frac{\partial}{\partial o_1} r_1 + 2 \sum \frac{\partial}{\partial q_1} s_1 + \frac{\partial}{\partial o} r \right)^j I \\ &= \sum_{j=0}^{\infty} \frac{\lambda^j}{j!} \left(\sum \frac{\partial}{\partial a_{m-k-l-l}} (k+1) a_{m-k-l-1l} + \frac{\partial}{\partial o} r \right)^j \Phi. \end{aligned} \quad (10)$$

When $o = o_1$, $r = r_1$, the first non-vanishing term of the right-hand series is the term where $j = \alpha$. Hence

THEOREM 3: *The α values of r constituting the set of r -coefficients of the α forms τ_i whose binary quadratic parts are factors of sum-multiplicity α of $x^m \phi_{0y/x} = 0$, are the roots of the following equation of order α :*

$$r^\alpha \frac{\partial^\alpha \Phi_{-o_1}^{(m)}}{\partial o_1^\alpha} + \binom{\alpha}{1} r^{\alpha-1} \frac{\partial^{\alpha-1} \Delta_k \Phi_{-o_1}^{(m)}}{\partial o_1^{\alpha-1}} + \binom{\alpha}{2} r^{\alpha-2} \frac{\partial^{\alpha-2} \Delta_k^2 \Phi_{-o_1}^{(m)}}{\partial o_1^{\alpha-2}} + \dots + \binom{\alpha}{\alpha} \Delta_k^\alpha \Phi_{-o_1}^{(m)} = 0, \quad (11)$$

where

$$\Delta_k = \sum_k \left(a_{0m-k-1} \frac{\partial}{\partial a_{1m-k-1}} + a_{1m-k-2} \frac{\partial}{\partial a_{2m-k-2}} + \dots + a_{m-k-10} \frac{\partial}{\partial a_{m-k-0}} \right) (k+1).$$

The corresponding values of q are the roots of the α linear equations

$$q_i \frac{\partial \Psi_{-n_i}^{(m)}}{\partial n_i} + \Delta_0 \Psi_{-n_i}^{(m)} = 0 \quad (i = 1, 2, \dots, \alpha). \quad (12)$$

In the functions Φ , Ψ used here the coefficients $a_0, a_1, a_2, \dots, a_m$ of § 3 are replaced by $a_{0m}, a_{1m-1}, \dots, a_{m0}$, respectively.

Next let $r^2 + o_1 r + n_{a_1+a_2+\dots+a_h}$ be a factor of ϕ_{0r} of multiplicity α_h , and of product-multiplicity α_h ($h = 1, 2, \dots, k$), and let these factors be all of sum-multiplicity $\alpha = \alpha_1 + \alpha_2 + \dots + \alpha_k$. Then equations (8), § 3, hold ($\xi_1 = -o_1$), so that $n_{a_1+a_2+\dots+a_h}$ is a root of $\Psi_{-n}^{(m)}$ of multiplicity α_h^2 ($h = 1, 2, \dots, k$). Also $r^2 + o_1 r + n_{a_1+a_2+\dots+a_h}$, being of multiplicity α_h in ϕ_{0r} , will be of multiplicity $\alpha_h - \kappa$ in $\phi_{\kappa r}$ ($\kappa = 0, 1, \dots, \alpha_h - 1$). Its product-multiplicity will be the same and its sum-multiplicity will be

$$\sum_{h=1}^k (\alpha_h - \kappa) = \alpha - \kappa k.$$

Then o_1 is a root of $\Phi_{-o}^{(m-\kappa)}$ of multiplicity

$$\mu_\kappa = (\alpha_1 - \kappa)^2 + (\alpha_2 - \kappa)^2 + \dots + (\alpha_k - \kappa)^2,$$

whereas $n_{a_1+a_2+\dots+a_h}$ is a root of $\Psi_{-n}^{(m-\kappa)}$ of multiplicity $(\alpha_h - \kappa)^2$.

Now we have

$$\Psi_{-n} = \Psi_{-n}^{(m)}(\alpha) = J(n_i, o_i, \dots; n) = J,$$

and $J(n_i, o_i, \dots; n_j) \equiv 0$ ($j = \alpha_1 + \alpha_2 + \dots + \alpha_h$). In J let n_i, o_i, q_i, n be replaced by $n_i + \lambda q_i, o_i + \lambda r_i, q_i + 2\lambda s_i, n + \lambda q$ respectively, and we get

$$\begin{aligned} & \sum_{j=0}^{\infty} \frac{\lambda^j}{j!} \left(\sum \frac{\partial}{\partial n_i} q_i + \sum \frac{\partial}{\partial o_i} r_i + 2 \sum \frac{\partial}{\partial q_i} s_i + \frac{\partial}{\partial n} q \right)^j J \\ &= \sum_{j=0}^{\infty} \frac{\lambda^j}{j!} \left(\sum \frac{\partial}{\partial a_{m-k-l-1}} (k+1) a_{m-k-l-1} + \frac{\partial}{\partial n} q \right)^j \Psi_{-n}^{(m)}. \end{aligned} \quad (13)$$

The first term of this series which does not vanish identically when $n = n_j$ is the one where $j = \alpha_h^2$. We get a similar result for r by replacing Ψ_{-n} by Φ_{-o} , $\frac{\partial}{\partial n} q$ by $\frac{\partial}{\partial o} r$, and α_h^2 by $\mu_0 = \alpha_1^2 + \alpha_2^2 + \dots + \alpha_k^2$. Thus we have

THEOREM 4: *The complete determination of the quadratic factors τ_i of a form f_{3m} , of general multiplicity, involves the solution for the q - and r -coefficients, of the equations*

$$r^{\mu_0} \frac{\partial^{\mu_0} \Phi_{-o_1}^{(m)}}{\partial o_1^{\mu_0}} + \binom{\mu_0}{1} r^{\mu_0-1} \frac{\partial^{\mu_0-1} \Delta_k \Phi_{-o_1}^{(m)}}{\partial o_1^{\mu_0-1}} + \binom{\mu_0}{2} r^{\mu_0-2} \frac{\partial^{\mu_0-2} \Delta_k^2 \Phi_{-o_1}^{(m)}}{\partial o_1^{\mu_0-2}} + \dots + \binom{\mu_0}{\mu_0} \Delta_k^{\mu_0} \Phi_{-o_1}^{(m)} = 0, \quad (14)$$

$$q^{\alpha_h^2} \frac{\partial^{\alpha_h^2} \Psi_{-n_j}^{(m)}}{\partial n_j^{\alpha_h^2}} + \binom{\alpha_h^2}{1} q^{\alpha_h^2-1} \frac{\partial^{\alpha_h^2-1} \Delta_k \Psi_{-n_j}^{(m)}}{\partial n_j^{\alpha_h^2-1}} + \dots + \binom{\alpha_h^2}{\alpha_h^2} \Delta_k^{\alpha_h^2} \Psi_{-n_j}^{(m)} = 0 \quad (15)$$

$(h = 1, 2, \dots, k; \mu_0 = \alpha_1^2 + \alpha_2^2 + \dots + \alpha_k^2).$

§ 5. *Semi-Resultant and Discriminant Matrices of Forms Representing n -Lines.*

By a semi-resultant we shall mean a necessary and sufficient matrix condition that an m -line and an n -line, each represented by a ternary form, should have a 1-line in common. The name will probably justify itself inasmuch as a semi-resultant of two ternary forms will be seen to be the natural analogue of a binary resultant. It is, however, seminvariantive instead of invariantive.

The author has proved in another paper* that the general m -line is represented by the ternary form

$$f_{3m} = x_2^m \phi_0 x_1/x_2 + x_2^{m-1} x_3 \phi_1 x_1/x_2 + D^{-1} (-1)^{\frac{1}{2}m(m-1)} \sum_{j=2}^m x_3^j \sum_{i=0}^{m-j} \frac{\Delta_1^{m-j-i} \Delta_2^i R_m}{\binom{m-j-i}{i}} x_1^{m-j-i} x_2^i = 0, \quad (16)$$

where

$$x_2^{m-k} \phi_k x_1/x_2 = a_{k0} x_1^{m-k} + a_{k1} x_1^{m-k-1} x_2 + \dots + a_{km-k} x_2^{m-k} \quad (k = 0, 1),$$

R_m is the resultant of $x_2^m \phi_0 x_1/x_2$ and $x_2^{m-1} \phi_1 x_1/x_2$, D is the discriminant of $x_2^m \phi_0 x_1/x_2$, and

$$\Delta_1 = m a_{00} \frac{\partial}{\partial a_{10}} + (m-1) a_{01} \frac{\partial}{\partial a_{11}} + \dots + a_{0m-1} \frac{\partial}{\partial a_{1m-1}},$$

$$\Delta_2 = m a_{0m} \frac{\partial}{\partial a_{1m-1}} + (m-1) a_{0m-1} \frac{\partial}{\partial a_{1m-2}} + \dots + a_{01} \frac{\partial}{\partial a_{10}}.$$

This result is a consequence of Theorem 1, § 1, when $\alpha_i = 1$.

* (Note added June 10, 1912.) This paper has been published in *Transactions Amer. Mathematical Society*, Vol. XII, No. 3 (1911), p. 373.

It is therefore natural to expect that *any joint invariant of an m -line and an n -line will be expressible entirely in terms of the two binary forms $x_2^{m-k} \phi_k x_1/x_2$ and the analogous forms from the equation of the n -line.* That this is really the case also follows from the results in § 1. That is, we may write the m -line and n -line equations as follows:

$$\begin{aligned} f_{3m} &= x_2^m \phi_0 x_1/x_2 + x_2^{m-1} x_3 \phi_1 x_1/x_2 + \dots, \\ g_{3n} &= x_2^n \psi_0 x_1/x_2 + x_2^{n-1} x_3 \psi_1 x_1/x_2 + \dots \quad (n \leq m). \end{aligned}$$

Then we have (Theorem 1)

$$\begin{aligned} f_{3m} &= \prod_{i=1}^m \left(x_1 + r_i x_2 - \phi_{1-r_i} / \frac{\partial \phi_{0-r_i}}{\partial r_i} \right), \\ g_{3n} &= \prod_{j=1}^n \left(x_1 + r_j x_2 - \psi_{1-r_j} / \frac{\partial \psi_{0-r_j}}{\partial r_j} \right). \end{aligned}$$

It now follows without difficulty that the condition for a common 1-line is equivalent to the condition that the following three binary equations have a common root:

$$\left. \begin{aligned} \phi_{0x} &= 0, & \psi_{0x} &= 0, \\ X_x^{m+n-2} &= \phi_{1x} \frac{\partial \psi_{0x}}{\partial x} - \psi_{1x} \frac{\partial \phi_{0x}}{\partial x} = 0. \end{aligned} \right\} \quad (17)$$

This condition can be expressed by means of a matrix, by using Stuyvaert's generalization* of the dialytic eliminant. Stuyvaert has shown that a necessary and sufficient condition that three polynomials in x of respective orders λ, μ, ν , $\nu \leq \lambda + \mu - 1$, may have a common root, is that a certain matrix of $\lambda + \mu$ rows and $\lambda + \mu + 1$ columns should be of rank $\dagger \lambda + \mu - 1$. We may express the condition here desired in the form of such a matrix with

$$\lambda = m - 1, \quad \mu = n, \quad \nu = m + n - 2.$$

* Stuyvaert: *Cinq études de géométrie analytique* (1908), p. 60.

† Conditions more desirable than $J_2 = J_3 = 0$ in § 1, part II, of my former paper can be obtained by constructing from the elements of J_2, J_3 , a Stuyvaert matrix, $\lambda = 2, \mu = 2, \nu = 3$, *e. g.*,

$$M = \begin{vmatrix} a & 0 & a' & 0 & a_0 \\ b & a & b' & a' & a_1 \\ c & b & c' & b' & a_2 \\ 0 & c & 0 & c' & a_3 \end{vmatrix},$$

where the sets a, b, c , and a', b', c' are respectively the non-zero elements of the third rows of J_2 and J_3 . The necessary conditions $J_2 = J_3 = 0$ are also sufficient if $a_x^2 = 0$ has but one real root. But the rank of M being 3 is both a necessary and sufficient condition, free from any assumption.

Thus, suppose $m = 3$, $n = 2$. Then

$$\begin{aligned}\phi_{0x} &= a_{00}x^3 + a_{01}x^2 + a_{02}x + a_{03}, & \psi_{0x} &= b_{00}x^2 + b_{01}x + b_{02}, \\ X_x^3 &= X_0x^3 + X_1x^2 + X_2x + X_3,\end{aligned}$$

where

$$\begin{aligned}X_0 &= 2a_{10}b_{00} - 3a_{00}b_{10}, \\ X_1 &= a_{10}b_{01} - 3a_{00}b_{11} + 2a_{11}b_{00} - 2a_{01}b_{10}, \\ X_2 &= a_{11}b_{01} - 2a_{01}b_{11} + 2a_{12}b_{00} - a_{02}b_{10}, \\ X_3 &= a_{12}b_{01} - a_{02}b_{11}.\end{aligned}$$

Then the semi-resultant of f_{33} and g_{32} is*

$$\rho_{32} \equiv \begin{vmatrix} a_{01}b_{00} - a_{00}b_{01} & a_{02}b_{00} - a_{00}b_{02} & a_{03}b_{00} & 0 \\ 0 & a_{01}b_{00} - a_{00}b_{01} & a_{02}b_{00} - a_{00}b_{02} & a_{03}b_{00} \\ b_{00} & b_{01} & b_{02} & 0 \\ 0 & b_{00} & b_{01} & b_{02} \\ X_0 & X_1 & X_2 & X_3 \end{vmatrix}$$

THEOREM 5: *A necessary and sufficient condition† that f_{3m} , g_{3n} may have a common linear factor is that ρ_{mn} should be of rank $m + n - 2$.*

It is worthy of notice that the coefficients of ϕ_{1x} , ψ_{1x} occur only in the X_i , so that two fourth-order determinants of ρ_{32} are linear in these coefficients. Therefore two of the latter are expressible rationally in terms of the remaining coefficients in the matrix.

THEOREM 6: *In general, if an m -line and an n -line have a 1-line in common, there will be $2m - 1$ independent coefficients in f_{3m} and $2n - 1$ in g_{3n} . Moreover, the remaining $\frac{1}{2}(m^2 - m + 2)$ coefficients of the first form and $\frac{1}{2}(n^2 - n + 2)$ of the second are all rationally expressible in terms of the $2(m + n - 1)$ independent ones.*

The condition that a given m -line $f_{3m} = 0$ may have a double 1-line can be expressed as a Stuyvaert matrix condition that the three binary equations

$$\phi_{0x} = 0, \quad \phi'_{0x} = 0, \quad \nabla_x^{2m-4} = \phi_{1x}'^2 - 2\phi_{0x}'\phi_{2x} = 0 \ddagger \quad (18)$$

* The negative condition

$$\begin{vmatrix} a_{01}b_{00} - a_{00}b_{01} & a_{02}b_{00} - a_{00}b_{02} & a_{03}b_{00} \\ b_{00} & b_{01} & b_{02} \end{vmatrix} \neq 0$$

is to be understood.

† As Stuyvaert points out, a certain negative condition concerning non-vanishing first minors of a definite determinant of order $\mu + \nu$ of the matrix is to be understood (see ρ_{32} above).

‡ Theorem 1 with $a_1 = 2$, $a_2 = 1$, $m = 3$.

may have a common root. When $m = 3$, this matrix is

$$\delta_3 \equiv \begin{vmatrix} 3a_{00} & 2a_{01} & a_{02} \\ 6a_{00}a_{02} - 2a_{01}^2 & 9a_{00}a_{03} - a_{02}a_{01} & 0 \\ 0 & 6a_{00}a_{02} - 2a_{01}^2 & 9a_{00}a_{03} - a_{02}a_{01} \\ 4a_{10}^2 - 12a_{00}a_{20} & 4a_{10}a_{11} - 12a_{00}a_{21} - 4a_{01}a_{20} & a_{11}^2 - 4a_{01}a_{21} \end{vmatrix}$$

THEOREM 7: *A necessary and sufficient condition that the m -line $f_{3m} = 0$ may have a double 1-line is that the matrix δ_m should be of rank $2m - 4$. The form f_{3m} then has but $2m - 2$ independent coefficients and the remainder are all rationally expressible in terms of those of ϕ_{0x} , ϕ_{1x} .*

Inasmuch as δ_m gives the condition for a repeated linear factor of f_{3m} , it might be called the *ultra-discriminant*. Generalizations giving conditions for a common 1-line for more than two m -lines, and corresponding results for p -ary forms $p > 3$, can be made readily.

Moreover, the method given may be extended in the direction of finding a necessary and sufficient condition for a 1-line of multiplicity greater than 2 of an m -line. Thus from Theorem 1 with $\alpha_1 = 3$ the condition for a triple 1-line is expressible as a Stuyvaert matrix (generalized) condition that the following five binary equations should have a common root:

$$\left. \begin{aligned} \phi_{0x} &= 0, & \phi'_{0x} &= 0, & \phi''_{0x} &= 0, \\ \rho_x^{2m-6} &= \phi_{1x}^{\prime\prime 2} - 2\phi_{0x}^{\prime\prime\prime}\phi'_{2x} = 0, \\ \sigma_x^{2m-6} &= 2\phi_{2x}^{\prime 2} - 3\phi_{1x}^{\prime\prime}\phi_{3x} = 0. \end{aligned} \right\} \quad (19)$$

These are derived from the Hessian of (4) with $\alpha_1 = 3$.

§ 6. Resolution into Rational Partial Fractions.

The results developed in sections 1, 3, 4 give us general formulas for the resolution of an ordinary rational proper algebraical fraction into the various standard types of partial fractions. For evidently the problem of resolving $x_2^{m-1}\phi_{1x_1/x_2}/x_2^m\phi_{0x_1/x_2}$ into partial fractions with real (linear and quadratic) denominators is identical, in part, with that of factoring a factorable form

$$x_2^m\phi_{0x_1/x_2} + x_2^{m-1}\phi_{1x_1/x_2} + \dots$$

into its linear and quadratic real ternary factors (see § 3). We can consider

the general partial fraction theorem, therefore, as one interpretation of theorems 3, 4. A sufficiently general statement of it is embodied in the following:

THEOREM 8: Let $x_2^m \phi_{0\ x_1/x_2} = \prod_{i=1}^h (x_1 + r_i x_2) \prod_{j=1}^l (x_1^2 + \xi_j x_1 x_2 + \eta_j x_2^2)$, where $x_1 + r_i x_2$ is a single (not multiple) factor and $x_1^2 + \xi_j x_1 x_2 + \eta_j x_2^2$ is of multiplicity 1, sum-multiplicity α_j and product-multiplicity 1. Then we have the resolution

$$\frac{x_2^{m-1} \phi_{1\ x_1/x_2}}{x_2^m \phi_{0\ x_1/x_2}} = \sum_{i=1}^h \frac{v_i}{x_1 + r_i x_2} + \sum_{j=1}^l \frac{\lambda_j x_1 + \mu_j x_2}{x_1^2 + \xi_j x_1 x_2 + \eta_j x_2^2},$$

where

$$v_i = \frac{\Delta_k \phi_{0-r_i}}{\Delta \phi_{0-r_i}}, \quad \mu_j = -\Delta_k \Psi_{-\eta_j}^{(m)} / \Delta \Psi_{-\eta_j}^{(m)},$$

and the numbers λ_j corresponding to the α_j factors of sum-multiplicity α_j are the α_j roots of

$$\lambda_{\alpha_j} \frac{\partial^{\alpha_j} \Phi_{-\xi_j}^{(m)}}{\partial \xi_j^{\alpha_j}} + \binom{\alpha_j}{1} \lambda_{\alpha_j-1} \frac{\partial^{\alpha_j-1} \Delta_k \Phi_{-\xi_j}^{(m)}}{\partial \xi_j^{\alpha_j-1}} + \binom{\alpha_j}{2} \lambda_{\alpha_j-2} \frac{\partial^{\alpha_j-2} \Delta_k^2 \Phi_{-\xi_j}^{(m)}}{\partial \xi_j^{\alpha_j-2}} \\ + \dots + \binom{\alpha_j}{\alpha_j} \Delta_k^{\alpha_j} \Phi_{-\xi_j}^{(m)} = 0,$$

where

$$\Delta = \frac{\partial}{\partial r}, \quad \Delta_k = \sum_k \left(a_{0\ m-k-1} \frac{\partial}{\partial a_{1\ m-k-1}} + a_{1\ m-k-2} \frac{\partial}{\partial a_{2\ m-k-2}} \right. \\ \left. + \dots + a_{m-k-10} \frac{\partial}{\partial a_{m-k0}} \right) (k+1).$$

These methods may be extended to the case of rational p -ary fractions $p > 2$.

PHILADELPHIA, PA., July, 1910.